

HEMATOLOGY

** This course meets the 1 hr. Hematology requirement for Florida licensees. **

A REVIEW OF BENZENE AND THE HEMATOLOGICAL SYSTEM

AUTHOR: O. Ifeanyi, M.D.

COURSE CODE: H030
CONTACT HOURS: 1
COURSE LEVEL: Intermediate
FLORIDA CE BROKER #: 20-722512
*** ALL COURSES ARE REVIEWED EVERY 2 YEARS ***

Continuing Education Unlimited

6231 PGA Blvd / Suite 104, #306 / Palm Beach Gardens, FL 33418
888-423-8462 / General Fax: 561-775-4933 / Answer Sheet ONLY Fax: 561-775-4948 / www.4CEUINC.com
PROVIDER #s: Florida: 50-2256 | ASCLS P.A.C.E.: 511 **ACCREDITING AGENCY #:** California: 0001

COURSE OBJECTIVES

At the end of this course you will be able to:

- 1.) Briefly discuss Benzene, including the common routes of exposure.
- 2.) Discuss different hematological disease states and how they're possibly affected by Benzene.
- 3.) Discuss safety policies, levels of exposure, and risk assessment associated Benzene exposure.
- 4.) Discuss low level exposures to Benzene, recalling genetic markers and chromosomal changes that may occur.

RIGHTSHOLDER:

Author: Obeagu Emmanuel Ifeanyi, M.D.

Publication: Blood Cells - An Overview of Studies in Hematology

Copyright: 2012, reviewed 2018

PERMISSIONS

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/3.0/us/>), which permits unrestricted use, distribution, and reproduction in any medium, for any purpose, provided the original work is properly cited.

ARTICLE REPRODUCTION

- ✓ Article reproduction falls under the guidelines of the Creative Commons License 3.0. You are free to:
SHARE: Copy & redistribute the material in any medium or format provided the work is properly cited.
ADAPT: remix, transform, & build upon, for any purpose, even commercially.
- ✓ Access a copy of the original, unedited document at:
<https://www.intechopen.com/books/blood-cell-an-overview-of-studies-in-hematology/platelets>
- ✓ Unlimited quizzes may be photocopied for sharing purposes only.

CEUINC LICENSE INFO:

LABORATORY

CA Department of Health:	0001
Florida Board of Clinical Lab:	50-2256
ASCLS P.A.C.E.®	511

Our courses are accepted by: AMTIE, AMT, ASCP, CA, FL, LA, ND, NV, MT, RI, TN, WV

** If you do not see your organization, state, or licensing agency listed above it does not mean that the credits will be unacceptable. Most licensing bodies accept **P.A.C.E.** credits, so please check directly with them for acceptance of our courses. **

PHLEBOTOMY

Most licensing bodies accept **P.A.C.E.** credits. Please check with them directly for acceptance of our course credits.

OTHER MEDICAL DISCIPLINES

Many medical licensing bodies will accept credits issued by valid licensed providers of other disciplines. Please check directly with the state, agency, or organization that issued your license for acceptance of our credits.

** CEUINC is approved as a provider of continuing education programs in the clinical laboratory sciences by the ASCLS P.A.C.E.® Program. **



Continuing Education Unlimited

6231 PGA Blvd, Suite 104 / #306 / Palm Beach Gardens, FL 33418
Phone: 888-423-8462 / Fax: 561-775-4933 / Web: www.4CEUINC.com / Email: Lab@4ceuinc.com

Thanks for choosing 4CEUINC for your continuing education needs! We strive to offer you current, cost effective course material. If you have comments or suggestions, be sure to add them to your evaluation. If you like our courses pass them on to a coworker or friend.

READ BEFORE COMPLETING MATERIAL



Last Revised 04/14/19

GENERAL COURSE INFO:

- 1.) Check your courses to make sure they are the correct titles that you ordered.
- 2.) All quiz answers can be found within the reading material and typically follow in order of the reading.
- 3.) All courses are reviewed every 2 yrs, at which time they will either be expired from our course list, updated, or renewed for an additional 2 years. Expiration dates can be found on this course list: https://4ceuinc.com/site/assets/pdf/current_course_list.pdf
- 4.) All of your records are available to you online through our website, even for Home Study. If you don't have a login to access your records, please contact our office and we will give you that information. **Please do not create a 2nd profile!**

HOME STUDY COURSES: (Combo Courses only)

- 1.) If you have multiple answer sheets, please make sure you're using the correct one.
- 2.) **CHANGES:** Home Study Complete formats have been replaced by "Book Only" formats, where you can purchase the printed book, then add on the quiz format option of your choice - - "Online Quiz Only" or "Answer Sheet Only".
- 3.) **CHANGES:** Answer sheets will no longer be mailed. You will print them out by logging into your online profile and visiting the My Courses list. Locate the correct course, then print your answer sheet. You may reprint as needed.
- 4.) **CHANGES:** As of May 1, 2019, answer sheets may **no longer** be returned by mail. You may submit them via fax to 561-775-4948 **OR** upload an image, doc, or pdf file by visiting your My Courses list online. As an alternate option, all Home Study quizzes **can be completed online**, which allows for **immediate** grading and certificate printing.
- 5.) **CHANGES:** As of May 1, 2019, we will no longer be returning Certificates of Completion. They may be printed out any time by visiting your "Course History" in your online profile. In the right column, take the Eval, then print the Certificate.

ONLINE COURSES:

- 1.) Course materials are currently offered in Adobe PDF documents, PowerPoint (PPT), and Video.
- 2.) You can read PDF & PPT documents online or print them out for reference. You may also store it on your hard drive or copy it to flash drive. Per copyright law, you're entitled to make **one copy** of each online course.
- 3.) Quizzes can be printed out if you'd like to work offline by clicking <Print Copy of Quiz>. Once you are done, simply transfer your answers to the active, online <Take Quiz> link and then click "Grade".
- 4.) When completing the online quiz, **it's important** to enter the answers without interruption, as some of the newer anti-virus programs may log you out of active content after a short period of time, causing you to lose your answers.

REGISTERED GROUPS:

- 1.) It's important to keep the Educational Coordinator's contact information current with our office at all times.
- 2.) In order to receive the group discount, you must be listed as a group member **OR** have the registration code to add to your online profile.
- 3.) You can share reading material with your co-workers for Combo Courses or Textbooks. They can purchase just the **online quiz** for that course allowing them to immediately complete the quiz and print their certificate of completion.
- 4.) **Have 5 or more employees?** Set up a Group Account & receive discounts on continuing education for your staff.



ALERT: When faxing, make sure to include a cover sheet with your full name & phone #. When faxing or uploading your Home Study answer sheets, make a note of the date and time you sent them! This safeguards you in the event that your answer sheet does not transmit properly to 4CEUINC.



Continuing Education Unlimited

6231 PGA Blvd, Suite 104 / #306 / Palm Beach Gardens, FL 33418
Phone: 888-423-8462 / Fax: 561-775-4933 / Web: www.4CEUINC.com / Email: Lab@4ceuinc.com



FREQUENTLY ASKED QUESTIONS

Last Revised 04/14/19

Q. Your phones are often busy, how can I reach you?

A. Because we have a small staff, the most efficient way would be to fill out a “Contact Us” form. On the main menu at www.4CEUINC.com, go to ‘Support’, ‘Contact Us’, then fill out the form. Be sure to enter your full information.

Q. I ordered a Home Study Combo Course, may I complete the quiz online?

A. Yes, quizzes from all course formats can be completed online, just log into your profile and go to “My Courses”.

Q. I need my certificate dated on a certain date; how can I be sure that this will happen?

A. 1.) Complete the quiz online. **OR** 2.) Be sure to fax or upload your answer sheet at least 72 hours before that date.

Q. What score is considered passing?

A. A score of 70% or higher is considered a passing grade. You may repeat the quiz if you fail it the first time.

Q. How can I return my Home Study answer sheet to your office?

A. CHANGES: As of May 15, 2019, answer sheets may no longer be returned by mail. They may **ONLY** be returned by fax to **561-775-4948** **OR** uploaded directly into our system through your My Courses list in your online profile. Be sure to keep a record of the date & time you send the Answer Sheet in the event that there is a transmission problem!

Q. May I share my Combo Course or Textbook with others?

A. YES, Combo Courses and Textbooks can be shared. One person will buy the “complete course” package & each additional person will purchase an “online quiz only”. Remember to share **both** the reading material **and** the quiz.

Q. How will I get my Certificate of Completion for a Home Study Combo Course?

A. CHANGES: As of May 1st, 2019, we will no longer be returning CEU certificates by mail due to past mail delivery problems. You may print your Certificate any time by logging into your online profile and visiting your course history.

Q. May I print out an online course?

A. YES, you may print **1 copy** of an online course. Copyright laws do not allow more than one copy to be printed!

Q. Where can I find additional information out about a course?

A. Basic course information is located on the course cover. Additional information is located on the 2nd page of each course.

Q. What are your most popular courses?

A. Currently our most popular courses remain the 12 Month Online Subscription and our Combo Courses.

Q. Does CEUINC offer group discounts or group packages?

A. YES. We require at least 5 participants and a person to act as the educational coordinator for the group. Discounted pricing depends on the number of participants and the package chosen. Please call 561-775-4944 Ext. 4 for details.

Q. Does CEUINC report to CE Broker for Florida licensees?

A. YES. Once you complete the online quiz, your course completions are sent to CE Broker immediately, provided that we have your correct Florida license number listed in your profile. Allow 48 hours for faxed or uploaded answer sheets.



ALERT: When faxing, make sure to include a cover sheet with your full name & phone #. When faxing or uploading your Home Study answer sheets, make a note of the date and time you sent them! This safeguards you in the event that your answer sheet does not transmit properly to 4CEUINC.



REVIEW OF BENZENE AND THE HEMATOLOGICAL SYSTEM

Categories: Hematology | Contact Hours: 1 | Course Code: H030

- 1.) Benzene is a chemical that is widely used in industrial settings, ranging from solvents to additives.
 - A. True
 - B. False

- 2.) There are ____ common routes of exposure to Benzene.
 - A. two
 - B. three
 - C. six

- 3.) Benzene has been known as a hematologic poison since the _____ century when aplastic anemia in workers fabricating tires was described.
 - A. eighteenth
 - B. nineteenth
 - C. twentieth

- 4.) Several case reports, case studies, and epidemiological studies demonstrate that MDS is caused by benzene exposure at benzene exposure levels less than 10ppm..
 - A. True
 - B. False

- 5.) In ____ OSHA reduced the permissible work place exposure from benzene from 10ppm to 1ppm, however, the US Supreme court invalidated that standard the following year.
 - A. 1949
 - B. 1978
 - C. 1992

- 6.) The use of chromosomal abnormalities as a biological marker of exposure in humans has become an important tool in the research.
 - A. True
 - B. False

******END OF QUIZ******

*****THIS PAGE INTENTIONALLY LEFT BLANK*****

A Review of Benzene and the Haematological System



Obeagu Emmanuel Ifeanyi*

Department of Health Services, Michael Okpara University of Agriculture, Nigeria

Submission: February 28, 2018; Published: March 20, 2018

*Corresponding author: Obeagu Emmanuel Ifeanyi, Department of Health Services, Michael Okpara University of Agriculture, Nigeria,

Summary

Benzene is a chemical substance that is widely used in industrial setting, ranging from solvent to gasoline additives. This shows that benzene is ubiquitous based on our daily activities. Although benzene is widely used in industrial setting, it has a toxic effect on the system of haematology, reproduction and neurology. The most probable mechanism of benzene-induced leukemia is through its phenolic metabolites acting in concert to produce DNA damage. This leads to mitotic recombination, chromosome translocations, and aneuploidy. These genotoxic events will, in turn, cause the activation of protooncogenes, loss of heterozygosity, and inactivation of tumor suppressor genes. In recent years, the relationship between benzene and smoking-induced hematopoietic malignancies has been solidified. Benzene has been known as a haematologic poison. Benzene has effect on all the haematopoietic cells and this happens by lodging in the bone marrow, leading to production of abnormal cells.

Keywords : Benzene; Hematology; Reproduction; Neurology; Leukemia

Abbreviations : IARC: International Agency for Research on Cancer's; MDS: Myelodysplastic Syndrome; NHL: Non-Hodgkin's Lymphoma; OSHA: Occupational Safety and Health Administration; EPA: Environmental Protection Agency; API: American Petroleum Institute; SCEs: Sister Chromatoid Exchanges; FISH: Fluorescence *In Situ* Hybridization; SP: Spectral Karyotyping; PCR: Polymerase Chain Reaction

Benzene

Benzene is reported as a chemical substance that is widely used in industrial settings, ranging from solvent to gasoline additives. Benzene is found in the air out of the emissions from cigarette smoke, motor vehicle exhaust, burning coal and oil, and gasoline service stations [1]. This shows that benzene is ubiquitous based on our daily activities. Although benzene is widely used in industrial settings, it has a toxic effect on the hematological, reproductive and neurological systems. Workers who are chronically overexposed can be inflicted with anemia, leucopenia, and thrombocytopenia. The peripheral blood morphology shows some immature leukocytes, such as bands, metamyelocytes, myelocytes, or myeloblasts [2]. Benzene is also a carcinogen in workers exposed to high levels of contaminated workplace air. It is shown that hundreds of leukemia cases caused by benzene exposure have been reported from some industries, such as the shoe industry, paint industry, and charcoal industry [3]. The occurrence of leukemia caused by benzene typically needs a long term exposure.

Similarly, epidemiological study shows that the leukemia cases increase gradually among the workers exposed to high levels of benzene. Benzene is one of the most studied causes of leukemia. The most probable mechanism of benzene-induced leukemia is through its phenolic metabolites acting in concert to produce DNA damage. This leads to mitotic recombination, chromosome translocations, and aneuploidy. These genotoxic events will, in turn, cause the activation of

protooncogenes, loss of heterozygosity, and inactivation of tumor suppressor genes. If this takes place in the bone marrow stem, a leukemic clone with selective growth advantage can arise. Epigenetic effects of benzene on the bone marrow stroma can then assist in the establishment of a leukemic clone. In the oil and natural gas mining industry, the employees have long term exposure to benzene, therefore, the possibility to get a carcinogenic effect from the benzene exposure is high [3].

Route of human exposure

Human exposure of significance is by the following routes:

- Inhalation
- Dermal exposure
- Ingestion of water & food contaminated with benzene

Although benzene is relatively soluble in water, commonly the magnitude of human exposure via water is probably negligible. The respiratory route is commonly the primary source of human exposure to benzene. Much of this exposure to the general population is by way of gasoline vapors and automobile emissions. In industrialized areas and heavily congested areas, levels of 15 parts per billion (ppb) up to 57ppb were described, while the average background levels have been reported to be 2.7 to 20ppb.

Benzene is absorbed through the skin, but skin contact is infrequent for the non-working general population. While the skin route is probably an insignificant source of exposure

for the general population, it has been shown as a significant route of exposure in the working population [3]. Smoking may be a significant benzene exposure source for a portion of the population. Studies have described levels of benzene exposure in active smokers at the range of 7.2 to 17.8ppb.

In recent years, the relationship between benzene and smoking-induced hematopoietic malignancies has been solidified. Korte et al. [4] combined epidemiological data on the health effects of smoking with risk assessment techniques for low-dose extrapolation and assessed the proportion of smoking-induced total leukemia and acute myeloid leukemia attributable to benzene and cigarette smoke. This study was based on linear potency models. According to this study, benzene is estimated to be responsible for approximately one-tenth to one-half of smoking-induced total leukemia mortality and up to three-fifths of smoking related acute myeloid leukemia mortality. Kasim et al. [4] has recently reported that active smoking was observed to be associated with a substantial increased risk of leukemia (odds ratio, OR = 1.5, 95% confidence interval, C.I. = 1.1 to 2.0). The International Agency for Research on Cancer's (IARC) 2004 monograph on cigarette smoking states that cigarette smoking causes leukemia, and that cigarettes contain sufficient quantity of the leukemogen benzene. This is an important development in terms of leukemogenesis and low-level exposure to benzene. One pack of cigarettes a day for 20 years is equivalent to 15ppb cumulative benzene exposure. This relatively new recognition further supports benzene leukemogenicity at levels lower than 1ppm and based on IARC, levels as low as 15ppb [5].

Benzene has been known as a haematologic poison since the nineteenth century when aplastic anemia in workers fabricating tires was described. Many other hematological diseases have since been reported to be the result of benzene exposure. Many of the hematological disorders related to benzene may not be dose-dependent as the mechanism of these diseases are yet not completely understood, although it is strongly believed that benzene carcinogenicity is mediated via immune suppression and DNA cell changes.

Myelodysplastic syndrome

Myelodysplastic Syndrome (MDS) is a bone marrow disease and is considered to be a preleukemic stage. Several case reports, case studies, and epidemiological studies demonstrate that MDS is caused by benzene exposure at benzene exposure levels less than 10ppm. It has been suggested that benzene-induced MDS is an early or predisposing event in the pathogenesis of benzene-induced hematologic diseases [6].

Aplastic anaemia/Pancytopenia

Aplastic anemia is a relatively rare, often fatal disorder. Its diagnosis is usually made on the basis of a significant reduction in the formed elements of the blood, including decreased white

blood cells, anemia, and thrombocytopenia. A decrease in all three of these blood cells counts is defined as pancytopenia. A marked decrease in the number of cells in the bone marrow is called aplastic anemia. It is accepted that these two are not two separate diseases but rather are part of a spectrum of bone marrow failure syndromes which can result from benzene toxicity. Indeed, a complete evaluation of a work force in a benzene-using plant revealed many affected individuals with effects ranging from a mild cytopenia to aplastic anemia of sufficient severity to warrant hospitalization; levels of exposure were 10-400ppm of benzene.

Acute myeloblastic leukemia

Acute myeloblastic leukemia is a cancer of the blood system in which there is an abnormal production of hematologic stem cells, granulocytic leukocytes, red blood cells and platelets. This disease is mostly observed in adults and has an increasing incidence with age, peaking in the 6th or 7th decade. There are a number of variants of acute myelogenous leukemia which can be considered to be part of the same disease. These include acute myelomonocytic leukemia, promyelocytic leukemia, and erythroleukemia.

The medical literature is replete with cases of acute myeloblastic leukemia in which benzene exposure has been shown as the causative agent. The relatively common description of aplastic anemia associated with benzene exposure followed through a pre-leukemic phase into acute leukemia further supports the concept that the bone marrow toxicity of benzene encompasses a wide spectrum of diseases presenting as anemia, thrombocytopenia, leukemia, or the other hematological diseases.

A published study in the New England Journal of Medicine, by Rinsky et al. quantitatively assessed the relationship between benzene exposure and leukemia, examining the mortality rate of cohorts with occupational exposure to benzene. Their findings are summarized below:

1. There's a strong positive exposure response relationship between benzene and leukemia.
2. On the basis of their study, they conclude that exposure levels of less than 1ppm annually, cumulative over a 40-year working lifetime increases the risk of leukemia by a factor of 1.7.
3. In the population studied, there was a statistically significant increase in death from multiple myeloma (MM is the cancer of plasma cells). Of interest in this study is a description of a patient who died from leukemia 34 years after his exposure to benzene levels of 19.56ppm over the years. Multiple myeloma, the cause of death in four members in this study, was described previously in relation to benzene, although in small numbers. Furthermore, it is of interest that these patients had a very long latency period from the time of exposure of over 20 years, This paper also demonstrates a latency as short as 1 year.

Lymphoma and lymphatic system

Recently, studies aimed at evaluating the effects of benzene and leukemia have also shown an increase in the relative risk of lymphatic system malignancies in benzene workers. A recent study by NIOSH described increased mortality from lymphoma and lymphocytic leukemia. A similar increased risk for lymphatic cancer has been reported by other investigators. Rubber chemical workers who were exposed to benzene had 4 to 5 fold higher risk of lymphoid malignancy than those unexposed [8].

Non-Hodgkin's lymphoma

Several lines of evidence demonstrate that benzene causes non-Hodgkin's lymphoma (NHL). Italian investigators presented a paper at the April 2005 American Association of Cancer Research. For benzene exposures of 15 years duration, researchers showed a significant excess risk for NHL and demonstrated a dose-response relationship when considering NHL subtypes. In 2004, Scandinavian researchers reported the results of a case-control study for NHL and occupational exposures. Risk of NHL was significantly increased for exposure to gasoline, oil products, and solvents [9].

Safety and policy

To reduce the risk of leukemia in industrial workers exposed to benzene, the United States Occupational Safety and Health Administration (OSHA) in 1978 reduced the permissible work place exposure of benzene from previous 10ppm to 1ppm. However, in 1980, the US Supreme court invalidated the OSHA benzene standard of 1ppm. The court states that OSHA had failed to provide substantial evidence of the need for regulation, and that it had not demonstrated a significant risk of material health impairment at the previous level of 10ppm. Since then, three studies have been published, in each of which the amount of benzene exposure has been found to correlate strongly with the risk of death from leukemia. The study published in the New England Journal of Medicine on benzene and leukemia further demonstrates that a cumulative benzene exposure of 400ppm is equivalent to a mean annual exposure of 10ppm over a 40 year working lifetime. They concluded that protection from benzene-induced leukemia would increase exponentially with any reduction in the permissible exposure limit enforceable to date. Obviously, the crucial question of who will develop a hematological disease as a result of exposure at the workplace to benzene is currently impossible to predict scientifically. Although a dose relation has been demonstrated, the fact that some cases have been described where exposure to benzene was not at excessive levels suggests that even strict protective efforts may not completely prevent industrially-related benzene exposure and hematological cancers [7].

Levels of exposure and risk assessment

The issue of what is a safe level of exposure to benzene and what is not a safe level of exposure, or as some would like to define it sufficient exposure to cause a hematological cancer has been

addressed by several studies and regulatory agencies. The study by the Environmental Protection Agency (EPA) as well as the International Agency for Research on Cancer [10] clearly indicates that there is no safe level of exposure to carcinogenic agents in the absence of epidemiological data of safety in humans. In the absence of safety studies in humans, experimental animal data must be applied from a policy and public health prevention point of view. Indeed, the American Petroleum Institute (API) stated that In as much as the body develops no tolerance to benzene and there is a wide variation in individual susceptibility, it is generally concluded that the only absolutely safe concentration for benzene is zero. Analysis of levels of exposure and risk assessment summarized by the 1998 position paper of the EPA clearly concludes that the dose-response relationship for benzene follows a linear line through zero.

The concept of cumulative benzene exposure for the working population must be well understood before one can address levels of exposure. They clearly show that diverse hematopoietic malignancies can develop at benzene exposure levels of less than 10ppm [11].

It is also important to remember that although many of the material safety data sheets of industrial solvents do not indicate the presence of benzene, the testimony in front of OSHA and the scientific papers published in that regards clearly indicate that industrial solvents contain benzene and cannot be produced without benzene contamination.

Benzene exposure levels are rarely available for most workers, because few workers are monitored for benzene exposure. The scientific medical literature allows the physician to extrapolate from the symptomatology of exposure, such as the threshold odor recognition for benzene, 61-91ppm, and symptomatology of dizziness, which is extrapolated to levels of 300ppm. This methodology has also been accepted by the U.S. Therefore it is imperative that the examining physician take a good history of exposure and look for odor recognition to extrapolate the levels of exposure and/or alternatively, symptoms of dizziness to extrapolate the levels of exposure. When history by the reporting physicians cannot be obtained, relying on depositions, job analysis and industrial hygienist assessment are acceptable [11].

Low Level Exposure To Benzene And Leukemia genetic studies and markers and chromosomal change

The use of chromosomal abnormalities as a biological marker of exposure in humans have become an important tool in the research. Several abnormalities are found, including structural and numerical chromosomal aberrations, sister chromatoid exchanges (SCEs) and micronuclear changes. These are markers of changes in the cellular genetic materials, and represent damage induced by chemicals. These methodologies are viewed as cytogenetic assays, and by themselves cannot provide a diagnosis, but they complement other methodologies which include gene mutation analysis, and DNA changes. Among the

important uses of cytogenetics as a biomarker is the relationship between chromosomal aberrations secondary to chemicals and carcinogenesis.

A patient who developed aplastic anemia after exposure to benzene, was shown to have significant chromatoid fragments. A cytogenetic study which was carried out later, on a patient who developed leukemia after 22 years of continuous exposure to a high concentration of benzene, showed that later in the process there were changes in chromosomes in the bone marrow [12].

While it is true that these findings are in agreement with previous studies [13] they still could not explain the 43% of the patients who were not exposed, and still had abnormal chromosomal changes. This is a very important observation, since some investigators in the field claim that the absence of chromosomal changes in benzene exposed individuals negates the clinical causative diagnosis of benzene induced hematopoietic disease. Essentially, all of the studies show that benzene can cause chromosomal changes, but does not cause perceptible changes in all patients, and the absence of visible chromosomal changes cannot and does not rule out the exposure to benzene as a causative factor. This is so because genetic point mutations and other types of changes are not observable using the standard cytogenetic banding techniques or even more sophisticated techniques such as fluorescence in situ hybridization (FISH), spectral karyotyping (SKY) or polymerase chain reaction (PCR) methods. Indeed, the courts have considered this issue and concluded that the genetic-chromosomal changes are not the equivalent of fingerprints of benzene exposure. The study by Zhang et al. [14] has examined chromosomal changes as a result of exposure to chemicals, such as benzene. Some leukemias have more typical chromosomal changes than others, but not all leukemias have typical chromosomal changes.

The study on the nature of chromosomal changes and benzene exposure was published by Luoping Zhang in *Critical Reviews in Toxicology*. The essence of this study is that benzene causes changes in certain chromosomes in certain hematological diseases. The absence of chromosomal changes does not rule out benzene exposure and benzene as a cause for the hematopoietic malignancy [14].

How to make or rule out a diagnosis of benzene-related hematological disorder

The examining physician who is faced with the question of causation in a patient with hematological malignancy and benzene exposure must utilize available epidemiologic and scientific data in the evaluation process. Ideally material safety data sheets as well as job descriptions and industrial hygiene reports, specifying the frequency and amount of exposure of benzene levels in the air should be considered, and the examining physician should request information in relation to other exposures such as solvent, radiation, pesticides, and cigarette smoking. Odour threshold recognition obtained via history, spills, headaches, and dizziness

are scientifically extrapolated to levels of exposure. Although the latency period may be important in the final analysis, one must remember that the scientific literature shows a range of anywhere from 6 months with an average of 12 years and up to 40 years in some cases. In some instances, it is probable that both the exposure to benzene on an industrial basis and exposure to other toxic chemical on a nonindustrial basis are additive to causation. In that scenario the reporting physician must determine whether the exposure to benzene, regardless of the other exposures, was a substantial factor in the development of the patient's hematological cancer [14].

Conclusion

Benzene is a hematological carcinogen based on both experimental animal studies and human studies, as well as in vitro studies. While the precise mechanism of benzene carcinogenicity is not clear evidence that benzene metabolites damage DNA and chromosomes, and in turn affects the stem cell: immature cells of the hematopoietic system which can in turn develop into any of the hematological cells originating from the bone marrow and the lymphatic system.

References

1. Ringen K, Addis P (1983) Protecting Workers From Benzene Exposure. In: Mehlman MA (Eds.), *Carcinogenicity and Toxicity of Benzene*, Princeton. Princeton Scientific Publishing pp. 77-89.
2. Holmberg B, Lundberg P (1985) Benzene Standards, Occurrence, and Exposure. *Am J Ind Med* 7(5-6): 373-383.
3. Kalnas J, Teitelbaum D (2000) Dermal Absorption of Benzene: Implications for Work Practices and Regulations. *Int J Occup Environ Health* 6(2): 114-121.
4. Korte JE, Hertz-Picciotto I, Schulz MR, Ball LM, Duell EJ (2000) The Contribution of Benzene to Smoking-Induced Leukemia. *Environ Health Perspect* 108(4): 333-339.
5. Kasim K, Levallois P, Abdous B, Auger P, Johnson KC (2005) Lifestyle factors and the risk of adult leukemia in Canada. *Cancer Causes Control* 16(5): 489-500.
6. Irons RD, Lv L, Gross SA, Ye X, Bao L, et al. (2005) Chronic exposure to benzene results in a unique form of dysplasia. *Leuk Res* 29(12): 1371-1380.
7. Rinsky RA (1987) Benzene and Leukemia: An Epidemiologic Risk Assessment. *Environ Health Perspect* 82: 189-191.
8. Wong O (1987) An Industry Wide Mortality Study of Chemical Workers Occupationally Exposed to Benzene. I. General Results. *Br J Ind Med* 44(6): 365-381.
9. Dryver E, Brandt L, Kauppinen T, Olsson H (2004) Occupational Exposures and Non-Hodgkin's Lymphoma in Southern Sweden. *Int J Occup Environ Health* 10(1): 13-21.
10. World Health Organization, International Agency for Research on Cancer (1987) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*. World Health Organization, Geneva, Switzerland, p. 1-42.
11. Brautbar N (2006) Leukaemia and low level benzene concentration: revisited. *European Journal of Oncology* 11(1).
12. Forni A, Moreo L (1967) Cytogenetic Studies in a Case of Benzene Leukaemia. *Eur J Cancer* 3(04): 251-255.

13. Golomb HM, Alimena G, Rowley JD, Vardiman JW, Testa JR, et al. (1982) Correlation of Occupation and Karyotype in Adults With Acute NonLymphocytic Leukemia. *Blood* 60(2): 404-411.
14. Zhang L, Eastmond DA, Smith MT (2002) The nature of chromosomal aberrations detected in humans exposed to benzene. *Crit Rev Toxicol* 32(1): 1-42.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/OABTJ.2018.02.555582](https://doi.org/10.19080/OABTJ.2018.02.555582)

*****THIS PAGE INTENTIONALLY LEFT BLANK*****